

JOURNAL OF FUNCTIONAL ANALYSIS **99**, 443–458 (1991)

Hodge Decomposition Along the Leaves of a Riemannian Foliation

JESÚS A. ALVAREZ LÓPEZ

*Department of Geometry and Topology, University of Santiago de Compostela,
15705 Santiago de Compostela, Spain*

AND

PHILIPPE TONDEUR

*Department of Mathematics, University of Illinois,
Urbana, Illinois 61801*

Communicated by R. B. Melrose

Received April 4, 1989; revised May 26, 1989

A Hodge decomposition theorem is proved for the leafwise Laplacian of a Riemannian foliation on a closed Riemannian manifold. © 1991 Academic Press, Inc.

1. INTRODUCTION AND RESULTS

Let \mathcal{F} be a smooth foliation on a closed oriented manifold M . Let $T\mathcal{F}$ be its tangent bundle and $Q = TM/T\mathcal{F}$, the normal bundle. Let g_M be a Riemannian metric on M , which decomposes TM as an orthogonal direct sum $T\mathcal{F} \oplus T\mathcal{F}^\perp$, and canonically identifies $T\mathcal{F}^\perp$ with Q . This gives rise to the bigrading of the algebra of smooth differential forms $\Omega = \Omega_M$

$$\Omega^{u,v} = \Gamma(\Lambda^v T\mathcal{F}^* \otimes \Lambda^u Q^*) \cong \Gamma \Lambda^v T\mathcal{F}^* \otimes_{C^\infty(M)} \Gamma \Lambda^u Q^*. \quad (1.1)$$

The exterior derivative d decomposes as $d = d_{0,1} + d_{1,0} + d_{2,-1}$, where the double subscripts indicate the bidegrees of the bihomogeneous components. The spectral sequence associated to the filtration

$$F^r \Omega = \bigoplus_{u \geq r} \Omega^{u,\cdot} \quad (1.2)$$

satisfies then canonically $(E_0, d_0) \cong (\Omega, d_{0,1})$ (see, e.g., [1, 11]). A natural idea is to try to find a relation between the cohomology $E_1 = H(\Omega, d_{0,1})$ and the harmonic forms associated to the leafwise Laplacian Δ_0 , the

Laplacian canonically associated to $d_{0,1}$. If $\delta_{0,-1}$ denotes the formal adjoint of $d_{0,1}$, and $D_0 = d_{0,1} + \delta_{0,-1}$ the “Dirac operator along the leaves,” then $\Delta_0 = D_0^2$. These operators are also closely related to the operators used in the study of the index problem in [4, 17], and in the study of \mathcal{F} -harmonic measures in [7]. In the spirit of Milgram and Rosenbloom [13], one has to study the heat flow associated to Δ_0 , and its long-time behavior.

In [14] a similar study has been carried out for the basic Laplacian Δ_B acting on the subcomplex $\Omega_B \subset \Omega_M$ of basic forms of a Riemannian foliation, thereby reproving the Hodge decomposition theorem of [6, 12]. This succeeded because of formal ellipticity properties of Δ_B .

In contrast, the operator Δ_0 is defined on all forms but is only “elliptic along the leaves.” A first step towards the desired decomposition theorem can be made as follows. First one constructs a chain of Hilbert space completions $\{H_{0,r}\}_{r \geq 0}$ of Ω_M with $H_{0,r} \supset H_{0,r+1}$, and compatible extensions $\bar{d}_{0,1,r}$, $\bar{\delta}_{0,-1,r}$, $\bar{D}_{0,r} = \bar{d}_{0,1,r} + \bar{\delta}_{0,-1,r}$, which define self-adjoint extensions $\bar{\Delta}_{0,r} = \bar{D}_{0,r}^2: \text{Dom}(\bar{\Delta}_{0,r}) \subset H_{0,r} \rightarrow H_{0,r}$ of $\Delta_0: \Omega_M \rightarrow \Omega_M$. These induce canonical continuous extensions $\bar{d}_{0,1,\infty}$, $\bar{\delta}_{0,-1,\infty}$, $\bar{D}_{0,\infty} = \bar{d}_{0,1,\infty} + \bar{\delta}_{0,-1,\infty}$ and finally $\bar{\Delta}_{0,\infty} = \bar{D}_{0,\infty}^2: H_{0,\infty} \rightarrow H_{0,\infty}$ to the Fréchet space

$$H_{0,\infty} = \bigcap_{r \geq 0} H_{0,r}. \quad (1.3)$$

The heat flows associated to the operators $\bar{\Delta}_{0,r}$ for $t \geq 0$ yield semi-groups of bounded operators

$$H_{0,r} \xrightarrow{e^{-t\bar{\Delta}_{0,r}}} H_{0,r}. \quad (1.4)$$

For $t \rightarrow \infty$, these give rise to the following Hodge decomposition theorem.

THEOREM A. *Let \mathcal{F} be a smooth foliation on a closed Riemannian manifold. Then the Laplacian Δ_0 along the leaves yields an orthogonal direct sum decomposition*

$$H_{0,\infty} \cong \ker \bar{\Delta}_{0,\infty} \oplus \overline{\text{im } \bar{\Delta}_{0,\infty}} \cong \ker \bar{\Delta}_{0,\infty} \oplus \overline{\text{im } \bar{d}_{0,1,\infty}} \oplus \overline{\text{im } \bar{\delta}_{0,-1,\infty}}.$$

An abstract analogous result is proved in Section 2 in a formal functional analytic setting. In Section 3 this is applied to the situation at hand.

In the case of an ordinary closed Riemannian manifold (viewed as a one leaf foliation, with $T\mathcal{F} = TM$ and $Q = 0$), the Laplacian in question is the ordinary Laplacian $\Delta = \Delta_M$. Then the scalar Bochner–Weitzenböck formula (see, e.g., [9]), implies that the usual Sobolev norms are equivalent with the operator defined norms given by (2.1) below, so that $H_{0,\infty}$ is the usual infinite Sobolev space H_∞ of Ω_M . From the Sobolev Lemma it follows that $H_\infty \cong \Omega_M$, so that the decomposition of Theorem A implies

the usual Hodge decomposition on forms. As noted by Roe, this construction and interpretation works equally well for the case of complete oriented Riemannian manifolds of bounded geometry, see [18].

A similar interpretation of $H_{0,\infty}$ can be given for Riemannian foliations. The corresponding result proved in Section 4 is as follows.

THEOREM B. *Let the situation be as in Theorem A, and assume \mathcal{F} to be Riemannian and the metric bundle-like. Then the space $H_{0,\infty}$ is the space of elements in $L^2(\Omega_M)$ defined by L^2 -Cauchy sequences $(\omega_m)_{m \in \mathbb{N}}$ of smooth forms $\omega_m \in \Omega_M$, such that any sequence of smooth forms obtained from $(\omega_m)_{m \in \mathbb{N}}$ by leafwise derivatives of any order of its local coefficient functions is L^2 -Cauchy.*

Therefore, $H_{0,\infty}$ is formed by the elements in $L^2(\Omega_M)$ whose leafwise derivatives of any order exist and are also in $L^2(\Omega_M)$.

Some comments on leafwise cohomologies resulting from these considerations are finally made in Section 5.

The authors thank I. D. Berg and F. Kamber for helpful comments.

2. AN ABSTRACT HODGE DECOMPOSITION THEOREM

We begin with an arbitrary complex Hilbert space H , and a symmetric (not necessarily bounded) operator S on H with dense domain $D \subset H$. Suppose that S verifies the hypothesis of Lemma 2.1 of Chernoff's paper [3], i.e., S maps D into itself, and there is a one-parameter group V_t of unitary operators on H such that $V_t D \subset D$, $V_t S = S V_t$ on D , and $(d/dt) V_t s = i S V_t s$ for $s \in D$. The definitions and results of this section are also valid for real Hilbert spaces, if the above conditions are satisfied by the corresponding complexifications.

Let $T = S^2$, which is a positive operator defined on D . For each integer $r \geq 0$, let $H_r(T)$ denote the Hilbert space completion of D with respect to the scalar product

$$\langle s, s' \rangle_r = \sum_{i=0}^r \langle T^i s, s' \rangle \quad \text{for } s, s' \in D. \quad (2.1)$$

For the corresponding norms $\| \cdot \|_r$, we have

$$r \leq r' \Rightarrow \|s\|_r \leq \|s\|_{r'} \quad \text{for all } s \in D. \quad (2.2)$$

Thus we obtain the chain of continuous inclusions

$$H = H_0(T) \supset H_1(T) \supset H_2(T) \supset \cdots \supset H_\infty(T), \quad (2.3)$$

where

$$H_\infty(T) = \bigcap_{r \geq 0} H_r(T) \quad (2.4)$$

is equipped with the obvious Fréchet topology.

Since the operator S is \langle, \rangle_r -symmetric, by [3, Lemma 2.1] each power of S is essentially self-adjoint in $H_r(T)$. Let \bar{T}_r be the closure of T in $H_r(T)$. By (2.2) we have

$$r \leq r' \Rightarrow \text{Dom}(\bar{T}_r) \supset \text{Dom}(\bar{T}_{r'}), \quad (2.5)$$

and the diagram

$$\begin{array}{ccc} \text{Dom}(\bar{T}_r) & \xrightarrow{\bar{T}_r} & H_r(T) \\ \cup & & \cup \\ \text{Dom}(\bar{T}_{r'}) & \xrightarrow{\bar{T}_{r'}} & H_{r'}(T) \end{array} \quad (2.6)$$

is commutative.

From definition (2.1) it follows that

$$H_{r+2}(T) \subset \text{Dom}(\bar{T}_r), \quad (2.7)$$

and the restriction

$$\bar{T}_r: H_{r+2}(T) \rightarrow H_r(T) \quad (2.8)$$

is a bounded operator. Moreover, the commutativity of (2.6) implies the commutativity of the diagram

$$\begin{array}{ccc} H_{r+2}(T) & \xrightarrow{\bar{T}_r} & H_r(T) \\ \cup & & \cup \\ H_{r+3}(T) & \xrightarrow{\bar{T}_{r+1}} & H_{r+1}(T). \end{array} \quad (2.9)$$

Hence the operators \bar{T}_r define a continuous operator

$$\bar{T}_\infty: H_\infty(T) \rightarrow H_\infty(T). \quad (2.10)$$

By the spectral theorem and the positivity of \bar{T}_r , we have the semi-group of bounded operators

$$e^{-t\bar{T}_r}: H_r(T) \rightarrow H_r(T) \quad \text{for } t \geq 0. \quad (2.11)$$

2.12. LEMMA. *The diagrams*

$$\begin{array}{ccc} H_r(T) & \xrightarrow{e^{-t\bar{T}_r}} & H_r(T) \\ \cup & & \cup \\ H_{r+1}(T) & \xrightarrow{e^{-t\bar{T}_{r+1}}} & H_{r+1}(T) \end{array}$$

are commutative.

Proof. Let \mathcal{B}_r be the subspace of the space of analytic vectors s (see [20], for example), such that for all $t \geq 0$ the series

$$\sum_{n=0}^{\infty} \frac{(-t\bar{T}_r)^n}{n!} s \quad (2.13)$$

is convergent in $H_r(T)$, and so its limit is $e^{-t\bar{T}_r}s$.

By (2.2) we have $\mathcal{B}_{r+1} \subset \mathcal{B}_r$. Then, by the commutativity of (2.6) and the density of \mathcal{B}_{r+1} in $H_{r+1}(T)$, the result follows. ▀

Since $e^{-t\bar{T}_r}$ converges strongly (as $t \rightarrow \infty$) to the orthogonal projection P'_0 of $H_r(T)$ onto the kernel of \bar{T}_r (by the spectral theorem), from (2.12) we obtain the commutativity of the diagram

$$\begin{array}{ccc} H_r(T) & \xrightarrow{P'_0} & H_r(T) \\ \cup & & \cup \\ H_{r+1}(T) & \xrightarrow{P'^{r+1}_0} & H_{r+1}(T). \end{array} \quad (2.14)$$

It follows that

$$\ker(\bar{T}_r) \supset \ker(\bar{T}_{r+1}). \quad (2.15)$$

From (2.12) we also have that the operators $e^{-t\bar{T}_r}$ define a continuous operator $e^{-t\bar{T}_\infty}: H_\infty(T) \rightarrow H_\infty(T)$.

For each $t > 0$

$$\psi_t(x) = \frac{e^{-tx} - 1}{x} \quad (2.16)$$

is a bounded continuous function on $[0, \infty)$, so we have the bounded operator $\psi_t(\bar{T}_r)$ on $H_r(T)$ defined by the spectral theorem. Using [5, p. 1199, Corollary 7], we obtain

$$\text{Dom}(\bar{T}_r \psi_t(\bar{T}_r)) = H_r(\bar{T}). \quad (2.17)$$

It follows that for all $s \in H_r(T)$

$$s = e^{-i\bar{T}_r} s - \bar{T}_r \psi_r(\bar{T}_r) s. \quad (2.18)$$

These are the same arguments as in [10, Lemma A.4]. It is immediate that

$$\ker \bar{T}_r \perp \operatorname{im} \bar{T}_r \quad \text{in } H_r(T). \quad (2.19)$$

Then (2.18) and (2.19) imply the following orthogonal direct sum decomposition

$$H_r(T) \cong \ker \bar{T}_r \oplus \overline{\operatorname{im} \bar{T}_r}. \quad (2.20)$$

From (2.5) and (2.6) we obtain further

$$\operatorname{im}(\bar{T}_r) \supset \operatorname{im}(\bar{T}_{r+1}). \quad (2.21)$$

Then, since $H_{r+1}(T)$ is $\|\cdot\|_r$ -dense in $H_r(T)$, from (2.15), (2.20), and (2.21) we obtain that $\ker(\bar{T}_r)$ and $\overline{\operatorname{im}(\bar{T}_r)}$ are the $\|\cdot\|_r$ -completions of $\ker(\bar{T}_{r+1})$ and $\operatorname{im}(\bar{T}_{r+1})$, respectively. But it is clear that on $\ker(\bar{T}_{r+1})$ we have $\|\cdot\|_{r+1} = \|\cdot\|_r$, and thus

$$\ker(\bar{T}_0) = \ker(\bar{T}_1) = \ker(\bar{T}_2) = \cdots = \ker(\bar{T}_\infty). \quad (2.22)$$

Therefore, since $\operatorname{im}(\bar{T}_\infty)$ is dense in each $\operatorname{im}(\bar{T}_r)$, the following fact results.

2.23. PROPOSITION. *Under the above conditions, we have the orthogonal direct sum decomposition*

$$H_\infty(T) \cong \ker(\bar{T}_\infty) \oplus \overline{\operatorname{im}(\bar{T}_\infty)}. \quad (2.24)$$

This result can be sharpened as follows.

2.25. PROPOSITION. *Under the same hypothesis, suppose that there is an integer $r \geq 0$ such that 0 is not an accumulation point of $\sigma(\bar{T}_r) - \{0\}$, where $\sigma(\bar{T}_r)$ is the spectrum of \bar{T}_r . Then we have orthogonal direct sum decompositions*

$$H_{r'}(T) \cong \ker(\bar{T}_{r'}) \oplus \operatorname{im}(\bar{T}_{r'}) \quad \text{for all } r' \geq r, \quad (2.26)$$

and

$$H_\infty(T) \cong \ker(\bar{T}_\infty) \oplus \operatorname{im}(\bar{T}_\infty). \quad (2.27)$$

Proof. By (2.5) and the commutativity of (2.6) we have

$$r \leq r' \Rightarrow \sigma(\bar{T}_r) \supset \sigma(\bar{T}_{r'}). \quad (2.28)$$

So, if the hypothesis is verified for some integer $r \geq 0$, it is also verified for any integer $r' \geq r$. Therefore, it is enough to prove (2.26) for $r' = r$.

Suppose that 0 is not an accumulation point of $\sigma(\bar{T}_r) - \{0\}$, and let $N = \inf(\sigma(\bar{T}_r) - \{0\}) > 0$. Then we have

$$\|\bar{T}_r s\|_r \geq N \cdot \|s\|_r \quad \text{for all } s \in \ker(\bar{T}_r)^\perp \cap \text{Dom}(\bar{T}_r). \quad (2.29)$$

Let \bar{T}_r^\perp be the restriction of \bar{T}_r to

$$\text{Dom}(\bar{T}_r^\perp) = \ker(\bar{T}_r)^\perp \cap \text{Dom}(\bar{T}_r). \quad (2.30)$$

\bar{T}_r^\perp is a self-adjoint operator on $\ker(\bar{T}_r)^\perp$. Then by (2.29) and the spectral theorem, we have that $1/\bar{T}_r^\perp$ is a well-defined bounded operator on $\ker(\bar{T}_r)^\perp$.

Again using [5, p. 1199, Corollary 7] we obtain

$$\text{Dom}(\bar{T}_r^\perp (1/\bar{T}_r^\perp)) = \ker(\bar{T}_r)^\perp, \quad (2.31)$$

and

$$\bar{T}_r^\perp \circ (1/\bar{T}_r^\perp) = \text{id}. \quad (2.32)$$

Therefore,

$$\ker(\bar{T}_r)^\perp = \text{im}(\bar{T}_r^\perp) = \text{im}(\bar{T}_r). \quad (2.33)$$

which implies (2.26).

To prove (2.27), it is enough to prove

$$\bigcap_{r' \geq r} \text{im}(\bar{T}_{r'}) = \text{im}(\bar{T}_\infty). \quad (2.34)$$

By assuming this we obtain that $\text{im}(\bar{T}_\infty)$ is closed in $H_\infty(T)$.

To prove (2.34) we observe that clearly $\text{im}(\bar{T}_\infty)$ is contained in each $\text{im}(\bar{T}_{r'})$. On the other hand, take $s \in H_\infty(T)$, so that for every $r' \geq r$ there exists $s_{r'} \in \text{Dom}(\bar{T}_{r'})$ such that $\bar{T}_{r'}(s_{r'}) = s$. Then we can suppose that $s_{r'} \in \ker(\bar{T}_{r'})^\perp$, obtaining that $s_{r'} = s_r$ for all $r' \geq r$ and $s \in \text{im}(\bar{T}_\infty)$. ■

Under the general hypothesis of this section we have the following lemmas.

2.35. LEMMA. *For all integers $r \geq 0$ we have*

$$\text{Dom}(\bar{T}_r) \subset H_{r+1}(T). \quad (2.36)$$

Proof. Let s be in $\text{Dom}(\bar{T}_r)$. Then there exists a sequence (s_n) in D

which is $\|\cdot\|_r$ -convergent to s and (Ts_n) is $\|\cdot\|_r$ -convergent in $H_r(T)$. For integers $n, m \geq 0$ we have then

$$\begin{aligned}\|s_n - s_{n+m}\|_{r+1}^2 &= \sum_{i=0}^{r+1} \langle T^i(s_n - s_{n+m}), s_n - s_{n+m} \rangle \\ &= \|s_n - s_{n+m}\|_0^2 + \langle Ts_n - Ts_{n+m}, s_n - s_{n+m} \rangle_r \\ &\leq \|s_n - s_{n+m}\|_r^2 + \|Ts_n - Ts_{n+m}\|_r \cdot \|s_n - s_{n+m}\|_r,\end{aligned}$$

and the last expression converges to zero as $n \rightarrow \infty$. Therefore, (s_n) is $\|\cdot\|_{r+1}$ -Cauchy, which implies that $s \in H_{r+1}(T)$. ■

2.37. LEMMA. For all $t \geq 0$ and all integers $r \geq 0$ we have

$$\text{im}(e^{-t\bar{T}_r}) \subset H_\infty(T). \quad (2.38)$$

Proof. Consider the real functions $f(x) = x$ and $g(x) = e^{-tx}$. Then by [5, p. 1199, Corollary 7] we have

$$\text{Dom}(f(\bar{T}_r)g(\bar{T}_r)) \subset \text{Dom}((fg)(\bar{T}_r)) \cap \text{Dom}(g(\bar{T}_r)) = H_r(T). \quad (2.39)$$

Hence, by (2.35) we obtain

$$\text{im}(e^{-t\bar{T}_r}) \subset \text{Dom}(\bar{T}_r) \subset H_{r+1}(T). \quad (2.40)$$

Therefore, by (2.12) we have

$$\text{im}((e^{-t\bar{T}_r})^k) \subset H_{r+k}(T) \quad \text{for all } k > 0, \quad (2.41)$$

and since

$$e^{-t\bar{T}_r} = (e^{-(t/k)\bar{T}_r})^k, \quad (2.42)$$

the result follows. ■

2.43. COROLLARY. The operators \bar{T}_r and \bar{T}_∞ have the same eigenvalues, and all their eigenvectors are in $H_\infty(T)$.

Proof. By the commutativity of (2.9) it is enough to prove that the eigenvectors of each operator \bar{T}_r are in $H_\infty(T)$. But if s is an eigenvector of \bar{T}_r and λ the corresponding eigenvalue, by the spectral theorem we have $e^{-\bar{T}_r s} = e^{-\lambda s}$, so $s \in H_\infty(T)$ by (2.37). ■

2.44. PROPOSITION. Assuming the conditions of (2.23), and in addition there is some $r \geq 0$ such that $H_r(T)$ is separable and the inclusion $H_{r+1}(T) \subset H_r(T)$ is compact, then there exists a complete orthonormal

system (COS) $\{s_i\}$ for $H_r(T)$, consisting of eigenvectors of \bar{T}_∞ in $H_\infty(T)$, and if $H_r(T)$ is of infinite dimension, then the corresponding eigenvalues satisfy $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \uparrow \infty$ (so they are of finite multiplicity).

Proof. We can suppose that $H_r(T)$ is of infinite dimension. As in the classical situation (see, e.g., [8]), one inverts the Friedrichs extension $T_{r,1}^F$ of the operator $\bar{T}_r + \text{id}$ to obtain a bounded operator

$$G_{r,1}^F = (T_{r,1}^F)^{-1}: H_r(T) \rightarrow D(\bar{T}_r) \cap H_{r+1}(T), \quad (2.45)$$

which composed with the compact inclusion into $H_r(T)$ yields a compact self-adjoint strictly positive operator on $H_r(T)$. Thus $G_{r,1}^F$ has positive eigenvalues $\mu_1 \geq \mu_2 \geq \dots \downarrow 0$ with corresponding eigenvectors $\{s_i\}$ constituting a COS for $H_r(T)$. Then it follows that $\bar{T}_r s_i = \lambda_i s_i$ with $\lambda_i = \mu_i^{-1} - 1$, satisfying $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \uparrow \infty$. Moreover, $s_i \in H_\infty(T)$ by (2.43). ■

Note that the hypothesis of (2.44) also implies the hypothesis of (2.25), so the orthogonal decompositions (2.26) and (2.27) also hold and \bar{T}_r is a Fredholm operator (with index zero, since it is self-adjoint).

Proposition 2.44 is the abstract result generalizing the situation of the usual Laplacian associated to a closed Riemannian manifold, where the compactness of the Sobolev inclusions is given by Rellich's Lemma. It also has the following converse.

2.46. PROPOSITION. *Under the hypothesis of (2.23), assume that H is separable and has a COS $\{s_i\}$ consisting of eigenvectors of T , such that the corresponding eigenvalues satisfy $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \uparrow \infty$. Then each inclusion $H_{r+1}(T) \hookrightarrow H_r(T)$ is compact.*

Proof. From (2.1) we obtain for any $r \geq 0$

$$\langle s, s' \rangle_r = \sum_{j=0}^r \langle \bar{T}_r^j s, s' \rangle \quad \text{for } s \in H_\infty(T) \text{ and } s' \in H_r(T). \quad (2.47)$$

Hence, by (2.43) each $H_r(T)$ has a COS $\{s_{r,i}\}$ given by

$$s_{r,i} = (\gamma_{r,i})^{-1/2} s_i, \quad \text{where } \gamma_{r,i} = \sum_{j=0}^r \lambda_i^j; \quad (2.48)$$

obtaining that

$$\langle s_{r,i}, s \rangle_r = \gamma_{r,i}^{1/2} \cdot \langle s_i, s \rangle \quad \text{for all } s \in H_r(T). \quad (2.49)$$

Let B be the unit ball in H_{r+1} . Given $\varepsilon > 0$ choose an integer $L > 0$ such that $L^{-1/2} < \varepsilon$. Since $0 \leq \lambda_1 \leq \lambda_2 \leq \dots \uparrow \infty$, there exists an integer $N > 0$

such that $\lambda_i \geq L$ for all $i \geq N$. Let Z be the space of elements $s \in H_{r+1}(T)$ so that $\langle s_i, s \rangle = 0$ for $i = 1, \dots, N$. Then for all $s \in B \cap Z$, we have by (2.48) and (2.49)

$$\begin{aligned} 1 > \|s\|_{r+1}^2 &= \sum_{i=1}^{\infty} \gamma_{r+1,i} \cdot |\langle s_i, s \rangle|^2 \\ &= \sum_{i=1}^{\infty} (\gamma_{r,i} \cdot \lambda_i + 1) \cdot |\langle s_i, s \rangle|^2 \\ &\geq L \cdot \|s\|_r^2, \end{aligned}$$

and thus $\|s\|_r < \varepsilon$. Now the argument follows as in the proof of Rellich's Lemma (see [19, Chapter III]). The unit ball of $H_{r+1}(T)/Z$ is precompact because Z is of finite codimension in $H_{r+1}(T)$, so it can be covered by finitely many balls of radius ε . Therefore, B can be covered by finitely many $\| \cdot \|_r$ -balls of radius 2ε , and so B is precompact in $H_r(T)$. ■

Let S_r be the closure of S in $H_r(T)$, which is self-adjoint. From the definition (2.1) it follows that

$$\text{Dom}(\bar{S}_r) = H_{r+1}(T), \quad (2.50)$$

and

$$\begin{array}{ccc} H_{r+1}(T) & \xrightarrow{\bar{S}_r} & H_r(T) \\ \cup & & \cup \\ H_{r+2}(T) & \xrightarrow{\bar{S}_{r+1}} & H_{r+1}(T) \end{array} \quad (2.51)$$

is a commutative diagram of bounded operators. Thus the operators \bar{S}_r define a continuous operator

$$\bar{S}_\infty : H_\infty(T) \rightarrow H_\infty(T). \quad (2.52)$$

It is easy to check that

$$\ker(\bar{S}_r) = \ker(\bar{T}_r) = \ker(\bar{T}_\infty) = \ker(\bar{S}_\infty), \quad (2.53)$$

and

$$\text{im}(\bar{S}_r) \perp \ker(\bar{S}_r), \quad \text{im}(\bar{S}_\infty) \perp \ker(\bar{S}_\infty). \quad (2.54)$$

Therefore, from (2.20) and (2.23) we obtain the orthogonal decompositions

$$H_r(T) \cong \ker(\bar{S}_r) \oplus \overline{\text{im}(\bar{S}_r)}, \quad (2.55)$$

and

$$H_{\infty}(T) \cong \ker(\bar{S}_{\infty}) \oplus \overline{\operatorname{im}(\bar{S}_{\infty})}. \quad (2.56)$$

Propositions similar to (2.25), (2.44), and (2.46) hold replacing \bar{T}_r and \bar{T}_{∞} by \bar{S}_r and \bar{S}_{∞} . But in this case there may be negative eigenvalues of \bar{S}_{∞} , so the results corresponding to (2.44) and (2.46) have to be stated in the appropriate way.

3. PROOF OF THEOREM A

We apply the results of the preceding section to a smooth foliation \mathcal{F} on a closed Riemannian manifold (M, g_M) . We have then the bigrading $\Omega^{u,v}$ of forms given by (1.2), and the decomposition $d = d_{0,1} + d_{1,0} + d_{2,-1}$. We can assume that M is oriented thus the formal adjoint δ of d can be given via the star operator $*$ on Ω (see, e.g., [15, Chap. IV]). It decomposes as $\delta = \delta_{0,-1} + \delta_{-1,0} + \delta_{-2,1}$ and for each (i, j) we have

$$\delta_{i,j} = (-1)^{n(r+1)+1} * d_{-i,-j} * \quad \text{on } \Omega^r, n = \dim M. \quad (3.1)$$

As explained in the Introduction, we then have the Dirac operator along the leaves

$$D_0 = d_{0,1} + \delta_{0,-1}, \quad (3.2)$$

and the Laplacian along the leaves

$$\Delta_0 = D_0^2 = d_{0,1} \delta_{0,-1} + \delta_{0,-1} d_{0,1}, \quad (3.3)$$

where the last equality follows from $d_{0,1}^2 = 0$, $\delta_{0,-1}^2 = 0$.

The complexification of the leafwise Dirac operator D_0 satisfies the hypothesis of Chernoff's Lemma 2.1 in [3]. This is obtained from [3, Corollary 1.4]. Then, with the notation of Section 2, we have the real Hilbert space $H_{0,r} = H_r(\Delta_0)$ and $H_{0,\infty} = H_{\infty}(\Delta_0)$. The operators D_0 and Δ_0 can be extended to continuous operators

$$\bar{D}_{0,\infty}, \bar{\Delta}_{0,\infty} : H_{0,\infty} \rightarrow H_{0,\infty}, \quad (3.4)$$

yielding the orthogonal decompositions

$$H_{0,\infty} \cong \ker \Delta_{0,\infty} \oplus \overline{\operatorname{im} \bar{\Delta}_{0,\infty}} \cong \ker \bar{D}_{0,\infty} \oplus \overline{\operatorname{im} \bar{D}_{0,\infty}}. \quad (3.5)$$

Moreover, the spaces $\Omega^{u,v}$ are $\langle, \rangle_{0,r}$ -orthogonal to each other for each

integer $r \geq 0$, where $\langle, \rangle_{0,r}$ denotes the scalar product of $H_{0,r}$. It follows that $\bar{D}_{0,\infty}$ can be decomposed as the sum of the continuous operators

$$\bar{d}_{0,1,\infty}, \bar{\delta}_{0,-1,\infty} : H_{0,\infty} \rightarrow H_{0,\infty}, \quad (3.6)$$

which are extensions of $d_{0,1}$ and $\delta_{0,-1}$, respectively. Since $\text{im } d_{0,1}$, and $\text{im } \delta_{0,-1}$ are $\langle, \rangle_{0,r}$ -orthogonal for each integer $r \geq 0$, from (3.5) we obtain the orthogonal decomposition

$$H_{0,\infty} \cong \ker \Delta_{0,\infty} \oplus \overline{\text{im } \bar{d}_{0,1,\infty}} \oplus \overline{\text{im } \bar{\delta}_{0,-1,\infty}}. \quad (3.7)$$

This is the Hodge decomposition along the leaves of \mathcal{F} stated in Theorem A.

4. PROOF OF THEOREM B.

In the same context as in the preceding section, assume now moreover that the foliation \mathcal{F} is Riemannian and the metric bundle-like. Let $p = \dim \mathcal{F}$, $q = \dim Q$, and $n = \dim M = p + q$.

We define a new connection on M by

$$\overset{\circ}{\nabla}_X Y = \pi_{\mathcal{F}} \nabla_X \pi_{\mathcal{F}} Y + \pi_Q \nabla_X \pi_Q Y \quad (4.1)$$

for vector fields X, Y , where $\nabla = \nabla^M$ is the Levi-Civita connection associated to g_M , and $\pi_{\mathcal{F}}, \pi_Q$ the orthogonal projections of TM onto $T\mathcal{F}, Q$, respectively. The induced connection on AT^*M verifies

$$\overset{\circ}{\nabla}_X = (\nabla_X)_{0,0} \quad \text{on } \Omega_M, \text{ for all vector fields } X. \quad (4.2)$$

From Reinhart's characterization of bundle-like metrics in [16], it follows that $\nabla_Y Y = 0$ for any infinitesimal transformation Y of \mathcal{F} of unit length, and orthogonal to \mathcal{F} . Thus $i_Y \nabla_Y \alpha = 0$ for all $\alpha \in \Omega^{0,\cdot}$. From this it is easy to obtain the following result using Koszul's formulas for d and δ (see, e.g., [15, Chap. IV]).

4.3. PROPOSITION. *Let E_1, \dots, E_p be a (local) orthonormal frame of $T\mathcal{F}$ on a distinguished chart $U \subset M$, and $\alpha_1, \dots, \alpha_p$ the dual co-frame of $T\mathcal{F}^*|_U$. Then for any $\omega \in \Omega^{0,\cdot}(U)$*

$$d_{0,1}\omega = \sum_{i=1}^p \alpha_i \wedge \overset{\circ}{\nabla}_{E_i} \omega \quad (4.4)$$

$$\delta_{0,-1}\omega = - \sum_{i=1}^p i_{E_i} \overset{\circ}{\nabla}_{E_i} \omega. \quad (4.5)$$

For the following we can assume without loss of generality that M and \mathcal{F} are oriented. Then the orientations of TM and $T\mathcal{F}$ induce an orientation of Q , yielding the star operators

$$*_\mathcal{F}: \Gamma A^v T^* \mathcal{F} \equiv \Omega^{0,v} \rightarrow \Omega^{0,p-v} \quad (4.6)$$

$$*_Q: \Gamma A^u Q^* \equiv \Omega^{u,0} \rightarrow \Omega^{q-u,0}, \quad (4.7)$$

with analogue properties to those of the usual star operator. A straightforward calculation gives the following lemma.

4.8. LEMMA. *We have according to the tensor expression (1.1)*

$$* = (-1)^{u(p-v)} *_\mathcal{F} \oplus *_Q: \Omega^{u,v} \rightarrow \Omega^{q-u,p-v}.$$

On a distinguished chart U we can find basic 1-forms $\beta_1, \dots, \beta_q \in \Omega^{1,0}(U)$, i.e., satisfying $d_{0,1} \beta_j = 0$, such that for each $x \in U$ the $(\beta_j)_x$ form an oriented orthonormal basis of Q_x^* . This implies the decomposition

$$\Omega^{u,v}(U) = \Omega^{0,v}(U) \otimes A^u \left(\bigoplus_{j=1}^q \mathbb{R} \cdot \beta_j \right). \quad (4.9)$$

4.10. PROPOSITION. *With respect to the decomposition (4.9), we have $D_0 = D_0 \otimes \text{id}$ on $\Omega(U)$.*

Proof. It is enough to prove $\delta_{0,-1} = \delta_{0,-1} \otimes \text{id}$, which is reduced to a simple sign verification by using (3.1) and (4.8). ■

With the same notation, each $\beta \in A(\bigoplus_{j=1}^q \mathbb{R} \cdot \beta_j)$ is supposed to be extended by zero to the whole manifold M . In this way, if $\beta \neq 0$ then it is not continuous on the boundary of U , but it defines an element in $L^2(\Omega_M)$.

4.11. LEMMA. *Let $\alpha, \alpha' \in \Omega_c(U)$ and $\beta, \beta' \in A(\bigoplus_{j=1}^q \mathbb{R} \cdot \beta_j)$. Then we have*

- (i) $\beta \wedge *_Q \beta' = \pm \langle \beta, \beta' \rangle (1/\text{Vol}(U))$. $\beta_1 \wedge \dots \wedge \beta_q$
- (ii) $\langle \alpha \wedge \beta, \alpha' \wedge \beta' \rangle = \pm \langle \alpha, \alpha' \rangle \langle \beta, \beta' \rangle (1/\text{Vol}(U))$.

Proof. We can assume β and β' to be of the same degree. Then we have $\beta \wedge *_Q \beta' \in \mathbb{R} \cdot \beta_1 \wedge \dots \wedge \beta_q$. Hence from (4.8) we obtain (i).

(ii) follows easily using (4.8) and (i). ■

Now, for each integer $r \geq 0$ let $\hat{\nabla}_\mathcal{F}^r: \Omega_M \rightarrow \Gamma(\otimes^r T\mathcal{F}^* \otimes AT^*M)$ be the operator defined by $\hat{\nabla}$ in the usual way,

$$(\hat{\nabla}_\mathcal{F}^r \omega)(X_1 \otimes \dots \otimes X_r) = \hat{\nabla}_{X_1} \dots \hat{\nabla}_{X_r} \omega \quad \text{for } X_i \in \Gamma T\mathcal{F}. \quad (4.12)$$

Also let further $\hat{\nabla}_\mathcal{F}^*$ denote the formal adjoint of $\hat{\nabla}_\mathcal{F} = \hat{\nabla}_\mathcal{F}^1$.

Using arguments similar to those in [19, Chap. 2], (4.3) yields the “leafwise Bochner–Weitzenböck formula,”

$$A_0 = \overset{\circ}{\nabla}^*_{\mathcal{F}} \overset{\circ}{\nabla}_{\mathcal{F}} + \overset{\circ}{R}_{\mathcal{F}} \quad \text{on } \Omega_M^{0,\cdot}, \quad (4.13)$$

where $\overset{\circ}{R}_{\mathcal{F}}$ is an operator of order zero involving the curvature of $\overset{\circ}{\nabla}$.

It can be seen easily that in general (4.13) does not hold on the entire Ω_M . Nevertheless, we have the following result.

4.14. LEMMA. *For each integer $r \geq 0$ there is a constant $c > 0$ such that if $\omega = \alpha \wedge \beta \in \Omega_c(U)$, with $\alpha \in \Omega_c^{0,\cdot}(U)$ and $\beta \in A(\bigoplus_{j=1}^q \mathbb{R} \cdot \beta_j)$, then we have*

- (i) $\|\omega\|_{0,r} \leq c \cdot \|\alpha\|_{0,r} \cdot \|\beta\|,$
- (ii) $\|\alpha\|_{0,r} \cdot \|\beta\| \leq c \cdot \|\omega\|_{0,r},$
- (iii) $\|\overset{\circ}{\nabla}_{\mathcal{F}}^r \omega\| \leq c \cdot \sum_{i=0}^r \|\overset{\circ}{\nabla}_{\mathcal{F}}^i \alpha\| \cdot \|\beta\|,$
- (iv) $\|\overset{\circ}{\nabla}_{\mathcal{F}}^r \alpha\| \cdot \|\beta\| \leq c \cdot \sum_{i=0}^r \|\overset{\circ}{\nabla}_{\mathcal{F}}^i \omega\|.$

Proof. (i) and (ii) follow directly from (4.10) and (4.13). For any integer $m > 0$ and for all vector fields X_1, \dots, X_m we have

$$\overset{\circ}{\nabla}_{X_1} \cdots \overset{\circ}{\nabla}_{X_m} \omega = (\overset{\circ}{\nabla}_{X_1} \cdots \overset{\circ}{\nabla}_{X_m} \alpha) \wedge \beta + \sum_{k=1}^{m-1} \sum_{\sigma} (\overset{\circ}{\nabla}_{X_{\sigma_1}} \cdots \overset{\circ}{\nabla}_{X_{\sigma_k}} \alpha) \wedge L_{\sigma} \beta, \quad (4.15)$$

where $\sigma = (\sigma_1, \dots, \sigma_k)$ ranges over the k -tuples of integers such that $1 \leq \sigma_1 < \sigma_2 < \dots < \sigma_k \leq m-1$, and L_{σ} is a linear map of $A(\bigoplus_{j=1}^q \mathbb{R} \cdot \beta_j)$ to $\Omega(U)$, which is defined using the connection $\overset{\circ}{\nabla}$. By taking a smaller open subset of U if needed, we can consider the image of each L_{σ} as a subspace of $L^2(\Omega_M)$. Then, since $A(\bigoplus_{j=1}^q \mathbb{R} \cdot \beta_j)$ is of finite dimension, the operators L_{σ} are bounded. Therefore, (iii) follows directly from (4.15).

The proof of (iv) follows easily by induction using (4.11)(ii) and (4.15). ■

4.16. PROPOSITION. *Each norm $\|\cdot\|_{0,r}$ is equivalent to the norm $\|\cdot\|'_{0,r}$ on Ω_M defined by*

$$\|\omega\|'_{0,r} = \left(\sum_{i=0}^r \|\overset{\circ}{\nabla}_{\mathcal{F}}^i \omega\|^2 \right)^{1/2}.$$

Proof. By (4.14) it is enough to prove that $\|\cdot\|_{0,r}$ is equivalent to $\|\cdot\|'_{0,r}$ on $\Omega^{0,\cdot}$, where it is a consequence of (4.13) by using a standard induction argument. ■

Now Theorem B follows directly from Proposition 4.16.

5. LEAFWISE COHOMOLOGIES

Under the hypothesis of Theorem A, $(H_{0,\infty}, \bar{d}_{0,1,\infty})$ is a topological differential space, and its cohomology, $H(H_{0,\infty}, \bar{d}_{0,1,\infty})$, has the induced topology. We define the reduced cohomology of $(H_{0,\infty}, \bar{d}_{0,1,\infty})$, denoted $\mathcal{H}(H_{0,\infty}, \bar{d}_{0,1,\infty})$ (or simply \mathcal{H}), to be the quotient of $H(H_{0,\infty}, \bar{d}_{0,1,\infty})$ by the closure of the trivial subspace. Then from Theorem A we obtain a canonical isomorphism

$$\ker(\bar{D}_{0,\infty}) \cong \mathcal{H}. \quad (5.1)$$

In the same way, the reduction \mathcal{E}_1 of E_1 is defined in [2] using the C^∞ -topology of Ω . There is a canonical homomorphism of \mathcal{E}_1 to \mathcal{H} , and hence also to $K = \ker(\bar{D}_{0,\infty})$.

The bigradation of Ω induces a bigradation on $H_{0,\infty}$, which defines a bigradation in both K and \mathcal{H} .

If M is oriented, by (3.1) the star operator $*$ induces isomorphisms

$$K^{u,v} \cong K^{q-u, p-v} \quad (5.2)$$

for all integers u and v . Thus, by (5.1) we also have duality isomorphisms

$$\mathcal{H}^{u,v} \cong \mathcal{H}^{q-u, p-v}. \quad (5.3)$$

If both M and \mathcal{F} are oriented, then according to the tensor decomposition (1.1) we have the operators $*_{\mathcal{F}} \otimes \text{id}$ and $\text{id} \otimes *_Q$ on Ω . Then, assuming \mathcal{F} to be Riemannian and the metric bundle-like, these operators also induce duality isomorphisms by (4.10)

$$K^{u,v} \cong K^{u, p-v}, \quad K^{u,v} \cong K^{q-u, v}, \quad (5.4)$$

$$\mathcal{H}^{u,v} \cong \mathcal{H}^{u, p-v}, \quad \mathcal{H}^{u,v} \cong \mathcal{H}^{q-u, v}. \quad (5.5)$$

Remark. In [17], J. Roe proves the following result for any smooth foliation \mathcal{F} on a closed Riemannian manifold M . If $\alpha \in \Gamma AT^* \mathcal{F} \equiv \Omega_M^{0,\cdot}$, then $e^{-t\bar{D}_{0,r}}\alpha \in \Omega_M^{0,\cdot}$ for any $t \geq 0$. When \mathcal{F} is Riemannian and the metric is bundle-like, by (4.10) we also have $e^{-t\bar{D}_{0,r}}(\Omega_M) \subset \Omega_M$. Considering the limit as $t \rightarrow \infty$, in general the orthogonal projection of $H_{0,\infty}$ on K does not preserve Ω_M . This can easily be seen for the Reeb foliation on S^3 . Nevertheless we conjecture that it is true when \mathcal{F} is Riemannian and the metric bundle-like, for in this case the distance between two leaves is locally constant, hence the limit as $t \rightarrow \infty$ of $e^{-t\bar{D}_{0,r}}\alpha$ should not produce

"transverse singularities" when α is smooth. If this conjecture were true, we would have the orthogonal decomposition

$$\Omega_M = \text{Ker}(D_0) \oplus \overline{\text{Im}(D_0)}, \quad (5.6)$$

and the canonical isomorphism

$$\mathcal{E}_1 \cong \text{Ker}(D_0). \quad (5.7)$$

REFERENCES

1. J. A. ALVAREZ LÓPEZ, A finiteness theorem for the spectral sequence of a Riemannian foliation, *Illinois J. Math.* **33** (1989), 79–92.
2. J. A. ALVAREZ LÓPEZ, Duality in the spectral sequence of Riemannian foliations, *Amer. J. Math.* **111** (1989), 905–926.
3. P. R. CHERNOFF, Essential self-adjointness of powers of generators of hyperbolic equations, *J. Funct. Anal.* **12** (1973), 410–414.
4. A. CONNES, Sur la théorie non-commutative de l'intégration, in "Lecture Notes in Math.," Vol. 725, pp. 19–143, Springer-Verlag, New York, 1979.
5. N. DUNFORD AND J. SCHWARTZ, "Linear Operators, Part II: Spectral Theory," Interscience, New York, 1963.
6. A. EL KACIMI AND G. HECTOR, Décomposition de Hodge basique pour un feuilletage riemannien, *Ann. Inst. Fourier* **36** (1986), 207–227.
7. L. GARNETT, Foliations, the ergodic theorem and Brownian motion, *J. Funct. Anal.* **51** (1983), 285–311.
8. P. GILKEY, Invariance theory, the heat equation, and the Atiyah–Singer index theorem, in "Mathematics Lecture Series," Vol. II, Publish or Perish, 1984.
9. M. GROMOV AND B. LAWSON, Positive scalar curvature and the Dirac operator on complete Riemannian manifolds, *Inst. Hautes Études Sci., Publ. Math.* **58** (1983), 295–408.
10. J. HEITSCH AND C. LAZAROV, A Lefschetz fixed point theorem for foliated manifolds, *Topology* **29** (1990), 127–162.
11. F. KAMBER AND PH. TONDEUR, Foliations and metrics, in "Progress in Math.," Vol. 32, pp. 103–152, Birkhäuser, Boston, MA, 1983.
12. F. KAMBER AND PH. TONDEUR, DeRham-Hodge theory for Riemannian foliations, *Math. Ann.* **277** (1987), 415–431.
13. A. N. MILGRAM AND P. C. ROSENBLUM, Harmonic forms and heat conduction. I. Closed Riemannian manifolds, *Proc. Nat. Acad. Sci.* **37** (1951), 180–184.
14. S. NISHIKAWA, M. RAMACHANDRAN, AND PH. TONDEUR, The heat equation for Riemannian foliations, *Trans. Amer. Math. Soc.* **319** (1990), 619–630.
15. W. A. POOR, "Differential Geometric Structures," McGraw-Hill, New York, 1981.
16. B. REINHART, Foliated manifolds with bundle-like metrics, *Ann. of Math.* **69** (1959), 119–132.
17. J. ROE, Finite propagation speed and Connes' foliation algebra, *Math. Proc. Cambridge Philos. Soc.* **102** (1987), 459–466.
18. J. ROE, An index theorem on open manifolds, I, *J. Differential Geom.* **27** (1988), 87–113.
19. J. ROE, Elliptic operators, topology and asymptotic methods, in "Pitman Res. Notes Math. Ser.," Vol. 179, Longman Sci. Tech., Harlow, 1988.
20. J. WEIDMAN, "Linear Operators on Hilbert Spaces," Springer-Verlag, New York, 1980.